Chronic Care Algorithm (A) Identify the factors which have delayed recovery and address them in the treatment plan Chronic MSD condition. Symptoms greater than (B) Active care measures include exercises and/or rehabilitation and education. Twelve weeks. (A) (C) Patient assessment indicates significant relief of pain and progress towards premorbid function. Most patients with mild cases after receiving an initial trial of care are discharged. Most patients with moderate cases that received prior care are discharges (D) Most cases reach MMI within 60-90 days of care. (E) Limitations in activities of daily living (ADL's)/function. (F) Refer to supportive criteria. Additional Treatment (up to 30 days) Focuses on: Passive Treatment not generally to exceed Prior treatment Yes No one visit per week Patient at MTB? trial for this Active Care measures to restore function episode? **Addressing Pain Behavior** (B) No Initial clinical trial of care not to exceed thirty days. Care to facilitate functional restoration. Passive Care not generally to exceed two times per week. (B) No Significant Discharge or Improvement? Referral. (C) Yes Based on treatment and Alter or progress with No scope of practice, has Maximum treatment plan (typically Clinical and Functional Improvement not to exceed 30 days) been met? (D) Yes No Significant Discharge/Elective Residuals? Care (E) Yes PRN/Supportive, Referral or Discharge (F)